

Regimen Reference Order – BRST – trastuzumab deruxtecan (ENHERTU)

ARIA: BRST – [trastuzumab deruxtecan (ENHERTU)]

Planned Course: Every 3 weeks until disease progression or unacceptable toxicity

Indication for Use: Breast Cancer Metastatic or Locally Advanced, HER2-Positive
OR

Breast Cancer Metastatic or Locally Advanced, HER2-Low

CVAD: At Provider's Discretion

Proceed with treatment if:

ANC equal to or greater than $1.5 \times 10^9/L$ AND Platelets equal to or greater than $50 \times 10^9/L$

❖ Contact Physician if parameters not met

SEQUENCE OF MEDICATION ADMINISTRATION

Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
Not Applicable		

Treatment Regimen – BRST – trastuzumab deruxtecan (ENHERTU)

Establish primary solution 500 mL of: D5W

Drug	Dose	CCMB Administration Guideline
dexamethasone	12 mg	Orally 30 minutes pre-chemotherapy
ondansetron	16 mg	Orally 30 minutes pre-chemotherapy
trastuzumab deruxtecan (ENHERTU®)	5.4 mg/kg	IV in D5W 100 mL <ul style="list-style-type: none"> Dose 1 to be infused over 90 minutes Dose 2 and subsequent to be infused over 30 minutes (if first dose well tolerated) Use 0.2 or 0.22 micron filter <i>* Pharmacy Alert: This is a look-alike and sound-alike medication. Refer to Additional Information</i> <i>*Alert: Ensure brand name on prescription label (indicated in brackets on prescription label) matches prescribed order</i>

All doses will be automatically rounded that fall within CCMB Approved Dose Bands. See Dose Banding document for more information

In the event of an infusion-related hypersensitivity reaction, refer to the 'Hypersensitivity Reaction Standing Order'

REQUIRED MONITORING

Cardiac Monitoring

- Left Ventricular Ejection Fraction (LVEF) monitoring is recommended at baseline and every 4 cycles for the first 2 years then every 4 to 8 cycles thereafter

All Cycles

- CBC, serum creatinine, urea, electrolytes, liver enzymes and total bilirubin as per Physician Orders

Cycle 1

- Full vital signs (temperature, heart rate, respiratory rate, blood pressure and O₂ saturation) at baseline and as clinically indicated
- Observe patient for 90 minutes after infusion (first dose). Full vital signs prior to discharge

Cycles 2 and 3 and Onwards

- Full vital signs (temperature, heart rate, respiratory rate, blood pressure and O₂ saturation) at baseline and as clinically indicated
- Observe patient for 30 minutes after infusion. Full vital signs prior to discharge

Cycles 4 and Onwards

- Full vital signs (temperature, heart rate, respiratory rate, blood pressure and O₂ saturation) at baseline and as clinically indicated
- No observation period required. Patient can be discharged from treatment room if stable whether they had a reaction or not

Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
dexamethasone	8 mg	Orally once daily on Days 2 and 3
metoclopramide	10 – 20 mg	Orally every 4 hours as needed for nausea and vomiting

DISCHARGE INSTRUCTIONS

- Patients should be instructed to contact their cancer team immediately if symptoms of hypersensitivity reactions occur after discharge
- Patient should report any new or worsening respiratory symptom such as dyspnea, cough or fever
- Reinforce applicable safe handling precautions of medications, blood and body fluids for 48 hours after completion of chemotherapy

ADDITIONAL INFORMATION

- trastuzumab deruxtecan (ENHERTU®) can cause interstitial lung disease and pneumonitis
- There is a risk of medication errors between trastuzumab deruxtecan (ENHERTU®), trastuzumab emtansine (KADCYLA®) and trastuzumab. In order to minimize the risk, check the regimen ordered, the vial labels, and the prescription label to ensure that the drug being prepared and administered is **trastuzumab deruxtecan (ENHERTU)**
- ARIA regimen **BRST – [trastuzumab deruxtecan (ENHERTU)]** is located in the “Metastatic/Advanced-HER2 Low” folder to order trastuzumab deruxtecan (ENHERTU) for HER-2 Low disease