

Regimen Reference Order - BRST – vinorelbine (weekly for 3 doses)

ARIA: BRST – [vinorelbine (weekly x 3) q28d]

Planned Course: Once a week for 3 weeks, then 1 week off until disease progression or unacceptable toxicity (1 cycle = 28 days)

Indication for Use: Breast Cancer Metastatic

CVAD: Preferred (VESICANT INVOLVED)

Proceed with treatment if:

ANC equal to or greater than $1 \times 10^9/L$ AND Platelets equal to or greater than $75 \times 10^9/L$

❖ Contact Physician if parameters not met

SEQUENCE OF MEDICATION ADMINISTRATION

Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
Not Applicable		

Treatment Regimen – BRST – vinorelbine (weekly for 3 doses)

Establish primary solution 500 mL of: normal saline

Drug	Dose	CCMB Administration Guideline
Days 1, 8 and 15		
vinorelbine	25 mg/m ²	IV in normal saline 50 mL over 6 to 10 minutes by gravity infusion <i>Slower or faster administration causes vein irritation</i>
normal saline	125 mL	IV over 15 minutes *Nursing Alert: This volume is to be administered after standard flush

In the event of an infusion-related hypersensitivity reaction, refer to the 'Hypersensitivity Reaction Standing Order'

REQUIRED MONITORING

All Cycles

Day 1

- CBC, serum creatinine, urea, liver enzymes and electrolytes as per Physician Orders

Days 8 and 15

- CBC

Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
metoclopramide	10 – 20 mg	Orally every 4 hours as needed for nausea and vomiting

DISCHARGE INSTRUCTIONS

- Instruct patient to continue taking anti-emetic(s) at home
- Reinforce applicable safe handling precautions of medications, blood and body fluids for 48 hours after completion of chemotherapy

ADDITIONAL INFORMATION

- Not applicable