Regimen Reference Order - CUTA - cemiplimab

ARIA: CUTA - [cemiplimab]

Planned Course: Every 21 days for up to 32 cycles

Indication for Use: Advanced Cutaneous Squamous Cell Carcinoma

Drug Alert: Immune Checkpoint Inhibitor

CVAD: At Provider's Discretion

Proceed with treatment if:

- ANC equal to or greater than 1.5 x $10^9/L$ AND Platelets equal to or greater than 50 x $10^9/L$
- AST/ALT equal to or less than 3 times the upper limit of normal
- Total bilirubin equal to or less than 1.5 times the upper limit of normal
- Creatinine clearance equal to or greater than 30 mL/minute
 - Contact Physician if parameters not met

SEQUENCE OF MEDICATION ADMINISTRATION

| | | Pre-treatment Requirements | | | |
|---|----------------|----------------------------|-------------------------------|--|--|
| ı | Drug | Dose | CCMB Administration Guideline | | |
| l | Not Applicable | | | | |

| Treatment Regimen – CUTA - cemiplimab | | | | |
|---|--------|---|--|--|
| Establish primary solution 500 mL of: normal saline | | | | |
| Drug | Dose | CCMB Administration Guideline | | |
| cemiplimab | 350 mg | IV in normal saline 100 mL over 30 minutes Use 0.2 or 0.22 micron filter | | |

In the event of an infusion-related hypersensitivity reaction, refer to the 'Hypersensitivity Reaction Standing Order'

REQUIRED MONITORING

All Cycles

- CBC, serum creatinine, urea, electrolytes, liver enzymes, total bilirubin, albumin, glucose, serum cortisol and TSH as per Physician Orders
- Medical oncologist or designate (i.e. family practitioner in oncology) must assess patient for immune -mediated adverse reactions prior to each cycle
- Full vital signs (temperature, heart rate, respiratory rate, blood pressure and O₂ saturation) at baseline and as clinically indicated
- No observation period is required. Patient can be discharged from treatment room if stable whether they had a reaction or not



ADULT CUTA - cemiplimab

| | | Recommended Support Medications | | | |
|---------------|------|---------------------------------|-------------------------------|--|--|
| l | Drug | Dose | CCMB Administration Guideline | | |
| None required | | | | | |

DISCHARGE INSTRUCTIONS

- Patient should be instructed to contact their cancer team immediately if symptoms of hypersensitivity reactions occur after discharge
- Confirm that patient has received the CCMB Immune Checkpoint Inhibitor Medical Alert wallet card
- Reinforce to patient the immune-mediated adverse reactions and importance of reporting immediately
 - For severe symptoms, the patient should be instructed to go to the nearest emergency room. Oncologist on call should be contacted

ADDITIONAL INFORMATION

cemiplimab is an Immune Checkpoint Inhibitor. Consult with oncologist for immune -mediated adverse reactions;
 corticosteroids are often indicated

