

Regimen Reference Order – GAST – DOCetaxel

ARIA: GAST – [DOCetaxel (gastric)]

Planned Course: Every 21 days until disease progression or unacceptable toxicity

Indication for Use: Gastroesophageal Cancer Metastatic

CVAD: At Provider’s Discretion

Proceed with treatment if:

ANC equal to or greater than 1.5 x 10⁹/L AND Platelets equal to or greater than 100 x 10⁹/L

❖ Contact Physician if parameters not met

SEQUENCE OF MEDICATION ADMINISTRATION

Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
dexamethasone	8 mg	Orally twice daily the day before DOCetaxel treatment and one dose the morning of DOCetaxel treatment (Self-administered at home)

Treatment Regimen – GAST – DOCetaxel

Establish primary solution 500 mL of: normal saline

Drug	Dose	CCMB Administration Guideline
DOCetaxel	75 mg/m ²	IV in normal saline 250 mL over 1 hour <i>Use non-DEHP bags and non-DEHP administration sets</i> <i>Concentration dependent drug: Pharmacy will adjust diluent volume to ensure drug stability</i>
normal saline	100 mL	ONLY for patients with a PORT IV over 12 minutes <i>*Nursing Alert: This volume is to be administered after standard flush</i>

In the event of an infusion-related hypersensitivity reaction, refer to the ‘Hypersensitivity Reaction Standing Order’

REQUIRED MONITORING

All Cycles

- CBC, biochemistry, AST, ALT, ALK and total bilirubin as per Physician Orders
- Full vital signs (temperature, heart rate, respiratory rate, blood pressure and O₂ saturation) at baseline and as clinically indicated
- No observation period required. Patient can be discharged from treatment room if stable whether they had a reaction or not

Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
metoclopramide	10 – 20 mg	Orally every 4 hours as needed for nausea and vomiting

DISCHARGE INSTRUCTIONS

- Patients should be instructed to contact their cancer team immediately if symptoms of hypersensitivity reactions occur after discharge
- Instruct patient to continue taking anti-emetic(s) at home
- Reinforce applicable safe handling precautions of medications, blood and body fluids for 48 hours after completion of chemotherapy

ADDITIONAL INFORMATION

- Not applicable