ADULT Updated: May 10, 2022

Regimen Reference Order – GAST - TOMOX

ARIA: GAST – [TOMOX]

Planned Course: Adjuvant: Every 21 days for 6-8 Cycles

OR

Metastatic: Until disease progression or unacceptable toxicity

Indication for Use: Colorectal Cancer Adjuvant OR Metastatic

CVAD: At Provider's Discretion

Proceed with treatment if:

ANC equal to or greater than 1.5 x $10^9/L$ AND Platelets equal to or greater than $100 \times 10^9/L$

Creatinine clearance greater than 65 mL/min

Contact Physician if parameters not met

SEQUENCE OF MEDICATION ADMINISTRATION

Pre-treatment Requirements				
Drug	Dose	CCMB Administration Guideline		
Not Applicable				

Treatment Regimen – GAST - TOMOX Establish primary solution 500 mL of: D5W				
Drug	Dose	CCMB Administration Guideline		
ondansetron	16 mg	Orally 30 minutes pre-chemotherapy		
dexamethasone	12 mg	Orally 30 minutes pre-chemotherapy		
raltitrexed	3 mg/m ²	IV in 50 mL D5W over 15 minutes		
oxaliplatin	130 mg/m ²	IV in 500 mL D5W over 2 hours		
•	matically rounded that fa	Il within the DSG Approved Dose Bands. See GAST DSG – Dose Banding		

In the event of an infusion-related hypersensitivity reaction, refer to the 'Hypersensitivity Reaction Standing Order'



ADULT GAST - TOMOX

REQUIRED MONITORING

All Cycles

• CBC and biochemistry (including renal function) as per Physician Orders

Recommended Support Medications					
Drug	Dose	CCMB Administration Guideline			
dexamethasone	8 mg	Orally once daily on Days 2 and 3			
metoclopramide	10 – 20 mg	Orally every 4 hours as needed for nausea and vomiting			

DISCHARGE INSTRUCTIONS

- Instruct patient to continue taking anti-emetic(s) at home
- Reinforce applicable safe handling precautions of medications, blood and body fluids for 48 hours after completion of chemotherapy

ADDITIONAL INFORMATION

- raltitrexed must be dose reduced if creatinine clearance is less than 65 mL/min
- · oxaliplatin causes cold intolerance and laryngopharyngeal dysesthesia
 - o no ice chips or cold drinks
- oxaliplatin may cause progressive, irreversible neuropathy
 - o dose modification may be required

