

Regimen Reference Order – GAST – de Gramont

ARIA: GAST - [de Gramont (ADJ)]

GAST - [de Gramont (MET)]

Planned Course: Adjuvant: Every 2 weeks for a total of 12 cycles

OR

Metastatic: Every 2 weeks until disease progression or unacceptable toxicity

Indication for Use: Colorectal Cancer Adjuvant OR Metastatic

CVAD: Required (Ambulatory Pump)

Proceed with treatment if:

ANC equal to or greater than $1.5 \times 10^9/L$ AND Platelets equal to or greater than $100 \times 10^9/L$

❖ Contact Physician if parameters not met

SEQUENCE OF MEDICATION ADMINISTRATION

Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
Not Applicable		

Treatment Regimen – GAST – de Gramont

Establish primary solution 500 mL of: normal saline

Drug	Dose	CCMB Administration Guideline
dexamethasone	8 mg	Orally 30 minutes pre-chemotherapy
leucovorin	400 mg/m ²	IV in D5W 500 mL over 90 minutes
fluorouracil	400 mg/m ²	IV Push over 5 minutes
fluorouracil	2400 mg/m ²	IV in D5W continuously over 46 hours by ambulatory infusion device

All doses will be automatically rounded that fall within CCMB Approved Dose Bands. See Dose Banding document for more information

In the event of an infusion-related hypersensitivity reaction, refer to the 'Hypersensitivity Reaction Standing Order'

REQUIRED MONITORING

All Cycles

- CBC, serum creatinine, urea, liver enzymes and total bilirubin as per Physician Orders

Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
metoclopramide	10 – 20 mg	Orally every 4 hours as needed for nausea and vomiting

DISCHARGE INSTRUCTIONS

- Instruct patient to continue taking anti-emetic(s) at home
- Ensure patient has received a home chemotherapy spill kit and instructions for use
- Reinforce applicable safe handling precautions of medications, blood and body fluids for 48 hours after completion of chemotherapy

ADDITIONAL INFORMATION

- Not applicable