

## Regimen Reference Order – GENU - gemcitabine

ARIA: GENU – [gemcitabine]

Planned Course: Every 28 Days for 6 cycles

Indication for Use: Urothelial Carcinoma

CVAD: At provider's discretion

### **Proceed with treatment if:**

**ANC equal to or greater than  $1.5 \times 10^9/L$  AND Platelets equal to or greater than  $100 \times 10^9/L$**

❖ Contact Physician if parameters not met

## SEQUENCE OF MEDICATION ADMINISTRATION

### Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
Not Applicable		

### Treatment Regimen – GENU - gemcitabine

Establish primary solution 500 mL of: normal saline

Drug	Dose	CCMB Administration Guideline
<b>Days 1, 8 and 15</b>		
dexamethasone	8 mg	Orally 30 minutes pre-chemotherapy
gemcitabine	1250 mg/m <sup>2</sup>	IV in normal saline 250 mL over 30 minutes

In the event of an infusion-related hypersensitivity reaction, refer to the 'Hypersensitivity Reaction Standing Order'

## REQUIRED MONITORING

All Cycles

- CBC, serum creatinine, liver enzymes and electrolytes as per Physician Orders

### Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
metoclopramide	10 – 20 mg	Orally every 4 hours as needed for nausea and vomiting

## DISCHARGE INSTRUCTIONS

- Instruct patient to continue taking anti-emetic(s) at home
- Reinforce applicable safe handling precautions of medications, blood and body fluids for 48 hours after completion of chemotherapy

---

**ADDITIONAL INFORMATION**

---

- Not applicable