

## Regimen Reference Order

### GYNE - CISplatin with Concomitant Radiation

ARIA: GYNE – [CISplatin-5-rad]

Planned Course: Weekly for 5 weeks with concomitant radiation

Indication for Use: Cervical Cancer

CVAD: At Provider's Discretion

**Proceed with treatment if:**

**Week 1**

- ANC equal to or greater than  $1.5 \times 10^9/L$  AND Platelets equal to or greater than  $75 \times 10^9/L$
- Creatinine clearance greater than 60 mL/min

**Week 2 and onwards**

- ANC equal to or greater than  $0.8 \times 10^9/L$  AND Platelets equal to or greater than  $75 \times 10^9/L$
- Creatinine clearance greater than 60 mL/min
- ❖ Contact Physician if parameters not met

#### SEQUENCE OF MEDICATION ADMINISTRATION

##### Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
Not Applicable		

##### Treatment Regimen – GYNE - CISplatin with Concomitant Radiation

Establish primary solution 500 mL of: normal saline

Drug	Dose	CCMB Administration Guideline
<b>Days 1, 8, 15, 22 and 29</b>		
aprepitant	125 mg	Orally 1 hour pre-chemotherapy
dexamethasone	12 mg	Orally 30 minutes pre-chemotherapy
ondansetron	16 mg	Orally 30 minutes pre-chemotherapy
OLANzapine	2.5 mg	Orally 30 minutes pre-chemotherapy
CISplatin	40 mg/m <sup>2</sup>	IV in normal saline 250 mL over 1 hour

Flush after each medication:

- 50 mL over 6 minutes (500 mL/hr)

In the event of an infusion-related hypersensitivity reaction, refer to the 'Hypersensitivity Reaction Standing Order'

## REQUIRED MONITORING

### Weekly

- CBC, biochemistry, serum creatinine, urea, liver enzymes and electrolytes as per Physician Orders

### Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
aprepitant	80 mg	Orally once daily on Days 2, 3, 9, 10, 16, 17, 23, 24, 30 and 31
dexamethasone	8 mg	Orally once daily on Days 2, 3, 4, 9, 10, 11, 16, 17, 18, 23, 24, 25, 30, 31 and 32
OLANzapine	2.5 mg	Orally the evening of Days 1, 8, 15, 22 and 29 and then twice daily on Days 2, 3, 4, 9, 10, 11, 16, 17, 18, 23, 24, 25, 30, 31 and 32. Also use OLANzapine 2.5 to 5 mg AS NEEDED for breakthrough nausea and vomiting (including Days 1 to 4, 8 to 11, 15 to 18, 22 to 25 and 29 to 32) up to a maximum of 10 mg per day. Contact clinic if nausea/vomiting is not adequately controlled

## DISCHARGE INSTRUCTIONS

- Instruct patient to continue taking anti-emetic(s) at home
- Reinforce applicable safe handling precautions of medications, blood and body fluids for 48 hours after completion of chemotherapy

## ADDITIONAL INFORMATION

- CISplatin is ototoxic and nephrotoxic
- CISplatin can cause hypomagnesemia
- Transfusions may be necessary in order to maintain a hemoglobin greater than 100 g/L
- If CISplatin needs to be held, radiation may continue
- Creatinine clearance is determined by using patient's actual body weight