

Regimen Reference Order – GYNE – gemcitabine + DOCEtaxel (MET)

ARIA: GYNE – [gemcitabine + DOCEtaxel (MET)]

Planned Course: Every 21 days until disease progression or unacceptable toxicity

Indication for Use: Uterine Leiomyosarcoma Metastatic

CVAD: At Provider's Discretion

Proceed with treatment if:

Cycle 1

ANC equal to or greater than $1.5 \times 10^9/L$ AND Platelets equal to or greater than $100 \times 10^9/L$

Cycle 2 and onwards

ANC equal to or greater than $1.2 \times 10^9/L$ AND Platelets equal to or greater than $75 \times 10^9/L$

❖ Contact Physician if parameters not met

SEQUENCE OF MEDICATION ADMINISTRATION

Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
Day 1		
Not Applicable		
Day 8		
dexamethasone	8 mg	Orally twice a day the day before DOCEtaxel treatment and one dose the morning of DOCEtaxel treatment (Self-administered at home)

Treatment Regimen – GYNE – gemcitabine + DOCEtaxel (MET)

Establish primary solution 500 mL of: normal saline

Drug	Dose	CCMB Administration Guideline
Day 1		
dexamethasone	8 mg	Orally 30 minutes pre-chemotherapy
gemcitabine	900 mg/m ²	IV in normal saline 250 mL over 90 minutes
Day 8		
metoclopramide	20 mg	Orally 30 minutes pre-chemotherapy
gemcitabine	900 mg/m ²	IV in normal saline 250 mL over 90 minutes
DOCEtaxel	100 mg/m ²	IV in normal saline 250 mL over 1 hour <i>Use non-DEHP bags and non-DEHP administration sets Concentration dependent drug: Pharmacy will adjust diluent volume to ensure drug stability</i>

normal saline	100 mL	ONLY for patients with a PORT IV over 12 minutes <i>*Nursing Alert: This volume is to be administered after standard flush</i>
---------------	--------	---

In the event of an infusion-related hypersensitivity reaction, refer to the 'Hypersensitivity Reaction Standing Order'

REQUIRED MONITORING

All Cycles

Days 1 and 8

- CBC, serum creatinine and liver enzymes as per Physician Orders

Day 8 Only

- Full vital signs (temperature, heart rate, respiratory rate, blood pressure and O₂ saturation) at baseline and as clinically indicated
- No observation period is required after DOCEtaxel administration. Patient can be discharged from treatment room if stable whether they had a reaction or not

Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
metoclopramide	10 – 20 mg	Orally every 4 hours as needed for nausea and vomiting

DISCHARGE INSTRUCTIONS

- Patients should be instructed to contact their cancer team immediately if symptoms of hypersensitivity reactions occur after discharge
- Instruct patient to continue taking anti-emetic(s) at home
- Reinforce applicable safe handling precautions of medications, blood and body fluids for 48 hours after completion of chemotherapy

ADDITIONAL INFORMATION

- Not applicable