

Regimen Reference Order – HEME – luspatercept (β-thalassemia)

ARIA: HEME – [luspatercept]

Planned Course: Every 21 days

Indication for Use: Transfusion-dependent anemia associated with β-thalassemia

CVAD: Not Required

Proceed with treatment if:

Hemoglobin less than 115 g/L

❖ Contact Hematologist if parameters not met

SEQUENCE OF MEDICATION ADMINISTRATION

Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
Not Applicable		

Treatment Regimen – HEME – luspatercept (β-thalassemia)

Drug	Dose	CCMB Administration Guideline
luspatercept	1 mg/kg*	Subcutaneous once every 3 weeks into the upper arm, thigh, and/or abdomen <i>*Alert: Maximum volume at each injection site is 1.2 mL</i>
<p><i>*Dose may be increased to 1.25 mg/kg at the hematologist’s discretion if patient does not achieve a reduction in RBC transfusion burden of at least a third from baseline after at least two consecutive doses (6 weeks) at 1 mg/kg starting dose</i></p> <p><i>*If hemoglobin increases by greater than 20 g/L within 3 weeks, in the absence of a transfusion, dose reduction may be required at the hematologist’s discretion</i></p>		

In the event of an infusion-related hypersensitivity reaction, refer to the ‘Hypersensitivity Reaction Standing Order’

REQUIRED MONITORING

All Cycles

- CBC, serum creatinine, urea, electrolytes and liver enzymes as per Physician Orders
- Blood pressure at each clinic visit
- Full vital signs (temperature, heart rate, respiratory rate, blood pressure and O₂ saturation) prior to each dose and as clinically indicated
- No observation period required. Patient can be discharged from treatment room if stable whether they had a reaction or not

Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
None required		

DISCHARGE INSTRUCTIONS

- Patients should be instructed to contact their cancer team immediately if symptoms of hypersensitivity reactions occur after discharge
- Confirm that females of childbearing potential received the manufacturer Patient Card (wallet card) at Cycle 1 by the clinic
- Instruct patient to report to clinic if they are experiencing injection site reactions that are bothersome

ADDITIONAL INFORMATION

- Patients with β -thalassemia may be at an increased risk of thromboembolic events while receiving treatment with luspatercept. Patients with a higher baseline risk of thromboembolic events (e.g. asplenic patients) may require thromboprophylaxis
- luspatercept may increase blood pressure
- luspatercept is teratogenic
- luspatercept is provided by an Access Program with requirements. Prescriber's Checklist must be completed and a Patient Card must be given to women of childbearing potential