

Regimen Reference Order

LYMP – R-GDP (Split Day SLOW for Cycle #1)

ARIA: LYMP – [R-GDP (Split Day SLOW on Cycle 1)]

Planned Course: Every 21 days up to 6 cycles
Indication for Use: Relapsed/Refractory Non-Hodgkin Lymphoma

CVAD: At Provider's Discretion

Proceed with treatment if:

Day 1

- ANC equal to or greater than $1 \times 10^9/L$ AND Platelets equal to or greater than $50 \times 10^9/L$
- Creatinine clearance greater than 45 mL/minute

Day 8

- Blood work not required to proceed with treatment
- ❖ Contact Hematologist if parameters not met

SEQUENCE OF MEDICATION ADMINISTRATION

Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
allopurinol*	300 mg	Orally once daily for 10 days to begin 3 days prior to Cycle 1 and at provider's discretion for subsequent cycles (Self-administered at home) *Only patients at risk of tumor lysis syndrome will be prescribed allopurinol

Treatment Regimen – R-GDP (Split Day SLOW for Cycle #1)

Establish primary solution 500 mL of: normal saline		
Drug	Dose	CCMB Administration Guideline
Cycle 1		
Day 1		
acetaminophen	650 mg	Orally 30 minutes prior to riTUXimab
cetirizine	10 mg	Orally 30 minutes prior to riTUXimab
dexamethasone	40 mg	IV in normal saline 50 mL over 15 minutes
Wait 30 minutes after completion of IV pre-medications before starting riTUXimab		
riTUXimab (IV brand name specific)	375 mg/m ²	Slow infusion: (if greater than 6 months since last riTUXimab dose or no previous riTUXimab): IV made up to a final concentration of 1 mg/mL in normal saline. Start at 50 mg/hr for 60 minutes and escalate infusion rate in 50 mg/hr increments every 30 minutes to a maximum of 400 mg/hr <i>*Nursing Alert: IV tubing is primed with riTUXimab</i>

		<p><i>*Alert: Ensure brand name on prescription label (indicated in brackets on prescription label) matches prescribed order</i></p> <p><i>*Alert: Pharmacy to ensure final volume on label</i></p>
OR		
		<p>Slow infusion: (if equal to or less than 6 months since last riTUXimab dose): IV made up to a final concentration of 1 mg/mL in normal saline. Start at 100 mg/hr for 30 minutes and escalate infusion rate in 100 mg/hr increments every 30 minutes to a maximum of 400 mg/hr</p> <p><i>*Nursing Alert: IV tubing is primed with riTUXimab</i></p> <p><i>*Alert: Ensure brand name on prescription label (indicated in brackets on prescription label) matches prescribed order</i></p> <p><i>*Alert: Pharmacy to ensure final volume on label</i></p>
Day 2		
magnesium sulfate	2 g	IV in normal saline 1000 mL over 2 hours (Pre hydration)
aprepitant	125 mg	Orally 1 hour pre-chemotherapy
ondansetron	16 mg	Orally 30 minutes pre-chemotherapy
dexamethasone	40 mg	Orally 30 minutes pre-chemotherapy
gemcitabine	1000 mg/m ²	IV in normal saline 250 mL over 30 minutes
CISplatin	75 mg/m ²	IV in normal saline 500 mL over 1 hour <i>*Alert: CISplatin infusion must be complete prior to mannitol administration</i>
mannitol	12.5 g	IV in normal saline 1000 mL over 2 hours (Post hydration)
Days 3 and 4		
dexamethasone	40 mg	Orally once daily in the morning with food (Self-administered at home)
Day 9		
dexamethasone	8 mg	Orally 30 minutes pre-chemotherapy
gemcitabine	1000 mg/m ²	IV in normal saline 250 mL over 30 minutes
Cycles 2 to 6		
Day 1		
acetaminophen	650 mg	Orally 30 minutes prior to riTUXimab
cetirizine	10 mg	Orally 30 minutes prior to riTUXimab
dexamethasone	40 mg	IV in normal saline 50 mL over 15 minutes
Wait 30 minutes after completion of IV pre-medications before starting riTUXimab		
riTUXimab (Subcutaneous)	1400 mg (1400 mg = 11.7 mL)	<p>Subcutaneous: Administer over 5 minutes into abdomen</p> <p>Syringe should be held in hand for 5 minutes to warm up and decrease viscosity</p> <p>Use 25G needle</p> <p><i>*Nursing Alert: Ensure subcutaneous riTUXimab formulation is</i></p>

		used (riTUXimab-hyaluronidase, human)
OR		
riTUXimab (IV brand name specific)	375 mg/m ²	<p>Rapid infusion: IV in normal saline over 90 minutes: Infuse 50 mL of a 250 mL bag (or 100 mL of a 500 mL bag) over 30 minutes, then infuse the remaining 200 mL (or 400 mL of a 500 mL bag) over 60 minutes</p> <p><i>*Alert: Ensure brand name on prescription label (indicated in brackets on prescription label) matches prescribed order</i></p> <p><i>*Alert: Pharmacy to ensure final volume on label</i></p> <p>Concentration dependent drug: Pharmacy will adjust diluent volume to ensure drug stability</p>
OR		
		<p>Slow Infusion: IV made up to a final concentration of 1 mg/mL in normal saline. Start at 100 mg/hr for 30 minutes and escalate infusion rate in 100 mg/hr increments every 30 minutes to a maximum of 400 mg/hr</p> <p><i>*Nursing Alert: IV tubing is primed with riTUXimab</i></p> <p><i>*Alert: Ensure brand name on prescription label (indicated in brackets on prescription label) matches prescribed order</i></p> <p><i>*Alert: Pharmacy to ensure final volume on label</i></p>
magnesium sulfate	2 g	IV in normal saline 1000 mL over 2 hours (Pre hydration)
aprepitant	125 mg	Orally 1 hour pre-chemotherapy
ondansetron	16 mg	Orally 30 minutes pre-chemotherapy
gemcitabine	1000 mg/m ²	IV in normal saline 250 mL over 30 minutes
CISplatin	75 mg/m ²	IV in normal saline 500 mL over 1 hour <i>*Alert: CISplatin infusion must be complete prior to mannitol administration</i>
mannitol	12.5 g	IV in normal saline 1000 mL over 2 hours (Post hydration)
Days 2, 3 and 4		
dexamethasone	40 mg	Orally once daily in the morning with food (Self-administered at home)
Day 8		
dexamethasone	8 mg	Orally 30 minutes pre-chemotherapy
gemcitabine	1000 mg/m ²	IV in normal saline 250 mL over 30 minutes
All doses will be automatically rounded that fall within the DSG Approved Dose Bands. See LYMP DSG – Dose Banding document for more information		

In the event of an infusion-related hypersensitivity reaction, refer to the 'Hypersensitivity Reaction Standing Order'

REQUIRED MONITORING

All Cycles

Day 1

- CBC, serum creatinine, urea, electrolytes, liver enzymes, LDH, total bilirubin, uric acid and albumin as per Physician Orders

Baseline blood pressure prior to magnesium infusion and repeat 15 minutes after start of magnesium infusion (day of CISplatin administration)

Day 8 (or Day 9 on Cycle 1)

- No blood work required

INTRAVENOUS riTUXimab

- Full vital signs (temperature, heart rate, respiratory rate, blood pressure and O₂ saturation) at baseline and as clinically indicated
- No observation period is required. Patient can be discharged from treatment room if stable whether they had a reaction or not

SUBCUTANEOUS riTUXimab

- Full vital signs (temperature, heart rate, respiratory rate, blood pressure and O₂ saturation) at baseline, at discharge and as clinically indicated
- **15 minute observation period required after each dose**

Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
Cycle 1		
aprepitant	80 mg	Orally once daily on Days 3 and 4
metoclopramide	10 – 20 mg	Orally every 4 hours as needed for nausea and vomiting
Cycles 2 to 6		
aprepitant	80 mg	Orally once daily on Days 2 and 3
metoclopramide	10 – 20 mg	Orally every 4 hours as needed for nausea and vomiting

DISCHARGE INSTRUCTIONS

- Patients should be instructed to contact their cancer team immediately if symptoms of hypersensitivity reactions occur after discharge
- Instruct patient to continue taking anti-emetic(s) at home
- dexamethasone is a cancer therapy in this treatment regimen. Remind patient to take dexamethasone at home
- Reinforce applicable safe handling precautions of medications, blood and body fluids for 48 hours after completion of chemotherapy

ADDITIONAL INFORMATION

- CISplatin is ototoxic and nephrotoxic
- CISplatin can cause hypomagnesemia
- Administering nurse must document any infusion-related reactions with any dose of riTUXimab
- Ensure there were **no Grade 3 or 4** infusion-related reactions with any previous dose prior to administering riTUXimab via subcutaneous injection or rapid infusion

ADULT

LYMP – R-GDP (Split Day SLOW for Cycle #1)

- Intravenous ritUXimab formulation is available from more than one manufacturer and uses several different brand names. Brand name will be indicated in brackets after ritUXimab. **Ensure prescription label matches the brand name on prescribed order for intravenous ritUXimab**