

Regimen Reference Order – LYMP - gemcitabine

ARIA: LYMP – [gemcitabine]

Planned Course: Every 28 Days for 3 to 6 cycles

Indication for Use: Mycosis Fungoides

CVAD: At provider's discretion

Proceed with treatment if:

Day 1

- ANC equal to or greater than $1 \times 10^9/L$ AND Platelets equal to or greater than $50 \times 10^9/L$

Days 8 and 15

- Blood work not required to proceed with treatment
- ❖ Contact Physician if parameters not met

SEQUENCE OF MEDICATION ADMINISTRATION

Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
Not Applicable		

Treatment Regimen – LYMP - gemcitabine

Establish primary solution 500 mL of: normal saline

Drug	Dose	CCMB Administration Guideline
dexamethasone	8 mg	Orally 30 minutes pre-chemotherapy on Days 1, 8 and 15
gemcitabine	1200 mg/m ²	IV in normal saline 250 mL over 30 minutes on Days 1, 8 and 15

In the event of an infusion-related hypersensitivity reaction, refer to the 'Hypersensitivity Reaction Standing Order'

REQUIRED MONITORING

All Cycles

Day 1

- CBC, serum creatinine, urea, liver enzymes and total bilirubin as per Physician Orders

Days 8 and 15

- No blood work required

Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
metoclopramide	10 – 20 mg	Orally every 4 hours as needed for nausea and vomiting

DISCHARGE INSTRUCTIONS

- Instruct patient to continue taking anti-emetic(s) at home
 - Reinforce applicable safe handling precautions of medications, blood and body fluids for 48 hours after completion of chemotherapy
-

ADDITIONAL INFORMATION

- Not applicable