

Regimen Reference Order – LYMP – oBINutuzumab + GDP

ARIA: LYMP – [oBINutuzumab + GDP]

Planned Course: Every 21 days for 6 cycles

Indication for Use: Non-Hodgkin Lymphoma

CVAD: At Provider's Discretion

Proceed with treatment if:

Day 1

- ANC equal to or greater than $1 \times 10^9/L$ AND Platelets equal to or greater than $50 \times 10^9/L$
- Creatinine clearance greater than 45 mL/minute

Day 8

- Blood work not required to proceed with treatment
- ❖ Contact Hematologist if parameters not met

SEQUENCE OF MEDICATION ADMINISTRATION

Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
allopurinol	300 mg	Orally once daily for 10 days to begin 3 days prior to Cycle 1 (Self-administered at home) Only patients at risk of tumor lysis syndrome will be prescribed allopurinol <u>Note:</u> allopurinol should not be prescribed beyond 10 days unless under the direction of the hematologist. See <i>Additional Information</i>

Treatment Regimen – LYMP – oBINutuzumab + GDP

Establish primary solution 500 mL of: normal saline

Drug	Dose	CCMB Administration Guideline
Cycle 1		
Day 1		
cetirizine	10 mg	Orally 30 minutes prior to oBINutuzumab
acetaminophen	650 mg	Orally 30 minutes prior to oBINutuzumab
dexamethasone	40 mg	IV in normal saline 50 mL over 15 minutes 1 hour prior to oBINutuzumab <i>*Nursing Alert: oBINutuzumab starts 1 hour after completion of dexamethasone infusion</i>

Wait 1 hour after completion of IV pre-medication(s) before starting oBINutuzumab

oBINutuzumab	100 mg	IV in normal saline 100 mL following administration rates below: <ul style="list-style-type: none"> • 0 to 60 minutes – 6 mL/hour • 60 to 120 minutes – 12 mL/hour • 120 minutes onwards – 24 mL/hour <p><i>*Alert: Pharmacy to ensure final volume in bag = 100 mL (1 mg/mL final concentration)</i></p> <p><i>*Nursing Alert: IV tubing is primed with oBINutuzumab</i></p>
Day 2		
cetirizine	10 mg	Orally 30 minutes prior to oBINutuzumab
acetaminophen	650 mg	Orally 30 minutes prior to oBINutuzumab
dexamethasone	40 mg	IV in normal saline 50 mL over 15 minutes 1 hour prior to oBINutuzumab <i>*Nursing Alert: oBINutuzumab starts 1 hour after completion of dexamethasone infusion</i>
Wait 1 hour after completion of IV pre-medication(s) before starting oBINutuzumab		
oBINutuzumab	900 mg	IV in normal saline 250 mL following administration rates below: <ul style="list-style-type: none"> • 0 to 30 minutes – 14 mL/hour • 30 to 60 minutes – 28 mL/hour • 60 to 90 minutes – 42 mL/hour • 90 to 120 minutes – 56 mL/hour • 120 to 150 minutes – 69 mL/hour • 150 to 180 minutes – 83 mL/hour • 180 to 210 minutes – 97 mL/hour • 210 to 240 minutes – 111 mL/hour <p><i>*Alert: Pharmacy to ensure final volume in bag = 250 mL (3.6 mg/mL final concentration)</i></p> <p><i>*Nursing Alert: IV tubing is primed with oBINutuzumab</i></p>
magnesium sulfate	2 g	IV in normal saline 1000 mL over 2 hours (Pre hydration)
aprepitant	125 mg	Orally 1 hour pre- chemotherapy
ondansetron	16 mg	Orally 30 minutes pre- chemotherapy
gemcitabine	1000 mg/m ²	IV in normal saline 250 mL over 30 minutes
CISplatin	75 mg/m ²	IV in normal saline 500 mL over 1 hour <i>*Alert: CISplatin infusion must be complete prior to mannitol administration</i>
mannitol	12.5 g	IV in normal saline 1000 mL over 2 hours (Post hydration)
Days 3 and 4		
dexamethasone	40 mg	Orally once daily in the morning with food (Self-administered at home)
Days 8		
cetirizine	10 mg	Orally 30 minutes prior to oBINutuzumab

acetaminophen	650 mg	Orally 30 minutes prior to oBINutuzumab
oBINutuzumab	1000 mg	<p>Slow Infusion: IV in normal saline 250 mL following administration rates below:</p> <ul style="list-style-type: none"> • 0 to 30 minutes – 25 mL/hour • 30 to 60 minutes – 50 mL/hour • 60 to 90 minutes – 75 mL/hour • 90 minutes onwards – 100 mL/hour <p><i>*Alert: Pharmacy to ensure final volume in bag = 250 mL (4 mg/mL final concentration)</i></p> <p><i>*Nursing Alert: IV tubing is primed with oBINutuzumab</i></p>
dexamethasone	8 mg	Orally 30 minutes pre-chemotherapy
gemcitabine	1000 mg/m ²	IV in normal saline 250 mL over 30 minutes
Day 15		
cetirizine	10 mg	Orally 30 minutes prior to oBINutuzumab
acetaminophen	650 mg	Orally 30 minutes prior to oBINutuzumab
oBINutuzumab	1000 mg	<p>Slow Infusion: IV in normal saline 250 mL following administration rates below:</p> <ul style="list-style-type: none"> • 0 to 30 minutes – 25 mL/hour • 30 to 60 minutes – 50 mL/hour • 60 to 90 minutes – 75 mL/hour • 90 minutes onwards – 100 mL/hour <p><i>*Alert: Pharmacy to ensure final volume in bag = 250 mL (4 mg/mL final concentration)</i></p> <p><i>*Nursing Alert: IV tubing is primed with oBINutuzumab</i></p>
Cycles 2 to 6		
Day 1		
dexamethasone	40 mg	Orally once in the morning with food (Self-administered at home)
cetirizine	10 mg	Orally 30 minutes prior to oBINutuzumab
acetaminophen	650 mg	Orally 30 minutes prior to oBINutuzumab
oBINutuzumab	1000 mg	<p>Rapid Infusion: IV in normal saline 250 mL following administration rates below:</p> <ul style="list-style-type: none"> • 0 to 30 minutes – 25 mL/hour • 30 to 93 minutes – 225 mL/hour <p><i>*Alert: Pharmacy to ensure final volume in bag = 250 mL (4 mg/mL final concentration)</i></p> <p><i>*Nursing Alert: IV tubing is primed with oBINutuzumab</i></p>
OR		

		<p>Slow Infusion: IV in normal saline 250 mL following administration rates below:</p> <ul style="list-style-type: none"> • 0 to 30 minutes – 25 mL/hour • 30 to 60 minutes – 50 mL/hour • 60 to 90 minutes – 75 mL/hour • 90 minutes onwards – 100 mL/hour <p><i>*Alert: Pharmacy to ensure final volume in bag = 250 mL (4 mg/mL final concentration)</i></p> <p><i>*Nursing Alert: IV tubing is primed with oBINutuzumab</i></p>
magnesium sulfate	2 g	IV in normal saline 1000 mL over 2 hours (Pre hydration)
aprepitant	125 mg	Orally 1 hour pre-chemotherapy
ondansetron	16 mg	Orally 30 minutes pre-chemotherapy
gemcitabine	1000 mg/m ²	IV in normal saline 250 mL over 30 minutes
CISplatin	75 mg/m ²	IV in normal saline 500 mL over 1 hour <i>*Alert: CISplatin infusion must be complete prior to mannitol administration</i>
mannitol	12.5 g	IV in normal saline 1000 mL over 2 hours (Post hydration)
Days 2, 3 and 4		
dexamethasone	40 mg	Orally once daily in the morning with food (Self-administered at home)
Day 8		
dexamethasone	8 mg	Orally 30 minutes pre-chemotherapy
gemcitabine	1000 mg/m ²	IV in normal saline 250 mL over 30 minutes

In the event of an infusion-related hypersensitivity reaction, refer to the 'Hypersensitivity Reaction Standing Order'

REQUIRED MONITORING

All Cycles

- CBC, serum creatinine, urea, electrolytes, liver enzymes, LDH, total bilirubin, uric acid and albumin as per Physician Orders
- Baseline blood pressure prior to magnesium infusion and repeat 15 minutes after start of magnesium infusion (day/s of CISplatin administration)

Day 8 (and Day 15 on Cycle 1)

- No blood work required

oBINutuzumab monitoring

- Full vital signs (temperature, heart rate, respiratory rate, blood pressure and O₂ saturation) at baseline and as clinically indicated
- No observation period is required after oBINutuzumab administration. Patient can be discharged from treatment room if stable whether they had a reaction or not

Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
Cycle 1		
aprepitant	80 mg	Orally once daily on Days 3 and 4
metoclopramide	10 – 20 mg	Orally every 4 hours as needed for nausea and vomiting
Cycles 2 to 6		
aprepitant	80 mg	Orally once daily on Days 2 and 3
metoclopramide	10 – 20 mg	Orally every 4 hours as needed for nausea and vomiting

DISCHARGE INSTRUCTIONS

- Patients should be instructed to contact their cancer team immediately if symptoms of hypersensitivity reactions occur after discharge
- Instruct patient to continue taking anti-emetic(s) at home
- dexamethasone is a cancer therapy in this treatment regimen. Remind patient to take dexamethasone at home
- Reinforce applicable safe handling precautions of medications, blood and body fluids for 48 hours after completion of chemotherapy

ADDITIONAL INFORMATION

- CISplatin is ototoxic and nephrotoxic
- CISplatin can cause hypomagnesemia
- The patient may be prescribed filgrastim. If filgrastim is started on cycle 1, filgrastim should start on Day 11 and should not be given on Day 15 (oBINutuzumab treatment day). The duration of filgrastim to be determined by the hematologist
- Administering nurse must document any infusion-related reactions with any dose of oBINutuzumab
- Ensure there were **no Grade 3 or 4** infusion-related reactions with the three preceding infusions prior to administering oBINutuzumab via rapid infusion. Patients will be switched to rapid infusion at Cycle 2, Day 1 if lymphocyte count is less than $5 \times 10^9/L$
- **Note: For Cycles 2 to 6**, an entry called **“Physician Reminder – oBINutuzumab infusion time 1 Units Insert Miscellaneous once”** will appear in the electronic drug order. No action is required. **This prompt is to remind the prescriber to confirm that patient is eligible for oBINutuzumab rapid infusion**
- For Cycle 1, Days 1 and 2, oBINutuzumab administration is 6 to 8 hours on average. Treatment should be booked for earliest morning appointment