

## Regimen Reference Order – LYMP – zanubrutinib

ARIA: LYMP – [zanubrutinib]

**Planned Course:** Once daily until disease progression or unacceptable toxicity (1 cycle = 30 days)

**Indication for Use:** Waldenström's Macroglobulinemia

**Proceed with treatment if:**

**ANC equal to or greater than  $0.75 \times 10^9/L$  AND Platelets equal to or greater than  $50 \times 10^9/L$**

❖ **Contact Hematologist if parameters not met**

### Treatment Regimen – LYMP – zanubrutinib

Drug	Dose	CCMB Administration Guideline
zanubrutinib	320 mg*	Orally once daily with or without food Swallow whole (Self-administered at home)

\*Alternate dosing schedule of 160 mg twice daily may be used at the prescriber's discretion

**zanubrutinib (Brukinsa®) available dosage strength: 80 mg capsule**

**Classification: Cytotoxic, Hazardous**

## REQUIRED MONITORING

### Baseline

- Hepatitis B serology
- EKG

### Throughout treatment

- CBC, serum creatinine, urea, electrolytes, liver enzymes, LDH, total bilirubin, uric acid and albumin as per Physician Orders
  - Every 2 to 4 weeks for the first 6 months, then
  - Every 12 weeks and as clinically indicated
- Monitor blood pressure and heart rate for tachycardia and irregular rhythm at every clinic visit
- At physician's discretion, EKG to assess for atrial fibrillation if tachycardia or irregular rhythm present

### Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
None required		

## INSTRUCTIONS FOR PATIENT

- Patients should report signs and symptoms of bleeding (i.e. excess bruising), palpitations, syncope and skin changes
- Patients should use necessary sun protection due to a potential increased risk of skin cancer
- When zanubrutinib is prescribed twice daily, doses of zanubrutinib should be separated by approximately 12 hours
- zanubrutinib has potential for drug-drug interactions. Patients should notify clinic prior to starting any new medication
- Avoid grapefruit and grapefruit juice, Seville oranges (i.e. orange marmalade) and starfruit
- Reinforce applicable safe handling precautions of medications, blood, and body fluids while on zanubrutinib

---

## ADDITIONAL INFORMATION

---

- Hematologist should be consulted regarding dosing of zanubrutinib pre- and post-surgery for patients undergoing a surgical procedure, including dental surgery, due to risk of bleeding regardless of platelet count